

JANUARY 2008
VOL. 53, NO. 1JOURNAL OF
CHEMICAL &
ENGINEERING
DATAЗаказ № 363 Пор. №. 91Журнал/книга имеет: 2 стр. _____

_____ табл. _____ карт.

№ № вып. 2008 v 53 n 1-2

Примечания:

Расписка отв. за подготовку в перепл.: p ll
3165The ACS Journal for Data
Issued Monthly

Vapor Pressures for 1,4-Dioxane + Tetrabutylammonium Nitrate, Water + Tetrabutylammonium Nitrate, and 1,4-Dioxane + Water + Tetrabutylammonium Nitrate

Alexander V. Kurzin,* Andrey N. Evdokimov, Victorija B. Antipina, Olesja S. Pavlova, and Vladimir E. Gusev

Faculty of Chemical Technology, Saint-Petersburg State Technological University of Plant Polymers (SPSTUPP-SPbGTURP), 4 Ivana Chernykh, Saint-Petersburg 198095, Russia

The vapor pressures of the 1,4-dioxane + tetrabutylammonium nitrate and water + tetrabutylammonium nitrate systems have been measured at different salt molalities. The vapor pressures of the 1,4-dioxane + water + tetrabutylammonium nitrate system have also been measured at 11 constant salt molalities (0.100, 0.250, 0.500, 0.750, 1.000, 1.500, 2.000, 2.500, 3.000, 3.500, and 4.000) mol·kg⁻¹. These systems have been studied at (288.15, 298.15, and 308.15) K with a modified Othmer still. The electrolyte nonrandom two-liquid model was used for the calculation of the activity coefficients of solvents in the binary systems.

Introduction

Organic salts are important and are usually used as intermediate chemicals, reaction catalysts, inhibitors to undesired reactions, supporting electrolytes, and surfactants. The electrolyte systems containing large organic ions (ammonium, borate, phosphonium, sulfonium, arsonium, hydrazinium, pyridinium, other organic salts, and ionic liquids) continue to represent an important area of theoretical interest as well.¹

Recently, the vapor pressures of the aqueous and nonaqueous solutions of several alkylammonium salts have been reported.^{2–10} In this study, the vapor pressures of the 1,4-dioxane + tetrabutylammonium nitrate (TBAN), water + TBAN, and 1,4-dioxane + water + TBAN systems were measured at different salt molalities at (288.15, 298.15, and 308.15) K with an Othmer-type equilibrium cell. No vapor pressure data for these systems were found in the literature, but there are many experimental data for the binary 1,4-dioxane + water system. The electrolyte nonrandom two-liquid (NRTL) model of Chen et al.¹¹ was used to calculate solvent activity coefficients in the binary systems.

Experimental Section

Materials. The 1,4-dioxane (*w* ≥ 99.9 %, Aldrich) was stored above 3 Å molecular sieves. Double-distilled and deionized water was used. Tetrabutylammonium nitrate (*w* ≥ 99.0 %, Fluka) was previously dried at 75 °C in a vacuum oven until a constant mass was reached.

Procedure. Mixtures consisting of 1,4-dioxane, water, and TBAN were prepared gravimetrically using an analytical balance (Ohaus Explorer Pro Balance) with an uncertainty of ± 0.1 mg. Known masses of salt were dissolved in 0.5 dm³ of solvent (or mixed solvent), and the solution was placed in the still. Then the system was closed and the still was heated until constant pressure and temperature were attained. The average uncertainty of the mole fraction is ± 0.005. For a fixed liquid-phase composition, at least three data points were taken of the total pressure at the target temperature of (288.15, 298.15, and 308.15) K. The uncertainty of the measured temperature was

Table 1. Vapor Pressures for Pure 1,4-Dioxane (1) and Water (2) from *T* = (288.15 to 308.15) K

<i>T</i> /K	<i>p</i> ₁ /kPa		<i>p</i> ₂ /kPa	
	lit. ^a	exptl. ^b	lit. ^a	exptl. ^b
288.15	2.691	2.703	1.669	1.706
298.15	4.986	4.975	3.166	3.172
308.15	8.129	8.119	5.547	5.556

^a Refs 16 and 17. ^b This work.




Figure 1. Modified Othmer's still for vapor pressure measurement: 1, distillation still; 2, steam jacket; 3, condenser; 4, condensed vapor receiver; 5, stopcock; 6, to manometer; 7, thermometer; 8, heating loop.

± 0.1 K. Vapor pressure data for pure water and 1,4-dioxane are presented in Table 1.

Apparatus. A modified static Othmer-type condensed vapor recirculating still¹² used for the present study is shown in Figure 1. The heating loop (8) ensures mixing and prevents bumping.

* To whom correspondence should be addressed. E-mail: zakora@mail.ru.

Table 2. Experimental and Literature Vapor Pressure Data of the Benzyltributylammonium Chloride (1) + Water (2) System from $T = (298.15 \text{ to } 303.15) \text{ K}$ as a Function of Salt Molality m_1

$m_1/(\text{mol} \cdot \text{kg}^{-1})$	$T/\text{K} = 298.15$		$T/\text{K} = 303.15$	
	$p_{\text{exptl}}/\text{kPa}^a$	$p_{\text{lit}}/\text{kPa}^b$	$p_{\text{exptl}}/\text{kPa}^a$	$p_{\text{lit}}/\text{kPa}^b$
0.1	3.153	3.14	4.205	4.19
0.5	3.116	3.12	4.153	4.15
1.0	3.091	3.09	4.110	4.10
1.5	3.074	3.08	4.071	4.08
2.0	3.032	3.03	4.006	4.01
3.0	2.971	2.95	3.854	3.87
3.5	2.893	2.91	3.807	3.81
4.0	2.820	2.83	3.762	3.77

^a This work. ^b Ref 3.**Table 3.** Activity Coefficients of 1,4-Dioxane γ_1 , Vapor Pressure p of the 1,4-Dioxane (1) + Tetrabutylammonium Nitrate (2) System from $T = (288.15 \text{ to } 308.15) \text{ K}$ as a Function of Salt Molality m_2

$m_2/(\text{mol} \cdot \text{kg}^{-1})$	$T/\text{K} = 288.15$		$T/\text{K} = 298.15$		$T/\text{K} = 308.15$	
	p/kPa	γ_1	p/kPa	γ_1	p/kPa	γ_1
0.100	2.907	1.0198	4.899	1.0212	7.987	1.0073
0.250	2.884	1.0112	4.854	1.0180	7.914	1.0014
0.500	2.853	1.0020	4.800	1.0104	7.802	0.9750
0.750	2.825	0.9814	4.751	1.0010	7.698	0.9610
1.000	2.796	0.9792	4.692	0.9814	7.601	0.9512
1.500	2.768	0.9670	4.639	0.9475	7.545	0.9370



Table 4. Activity Coefficients of Water γ_1 , Vapor Pressure p of the Water (1) + Tetrabutylammonium Nitrate (2) System from $T = (288.15 \text{ to } 308.15) \text{ K}$ as a Function of Salt Molality m_2

$m_2/(\text{mol} \cdot \text{kg}^{-1})$	$T/\text{K} = 288.15$		$T/\text{K} = 298.15$		$T/\text{K} = 308.15$	
	p/kPa	γ_1	p/kPa	γ_1	p/kPa	γ_1
0.100	1.691	1.0601	3.150	1.0203	5.607	1.0170
0.250	1.684	1.0560	3.135	1.0244	5.579	1.0110
0.500	1.674	1.0420	3.122	0.9918	5.534	0.9807
0.750	1.665	1.0307	3.100	0.9722	5.506	0.9630
1.000	1.657	1.0097	3.078	0.9643	5.472	0.9489
1.500	1.638	0.9707	3.043	0.9414	5.410	0.9211
2.000	1.621	0.9240	3.008	0.9291	5.348	0.9013
2.500	1.608	0.8935	2.983	0.9179	5.303	0.8892
3.000	1.593	0.8681	2.958	0.8908	5.258	0.8784
3.500	1.581	0.8492	2.939	0.8804	5.213	0.8760
4.000	1.565	0.8305	2.904	0.8713	5.157	0.8688

The electrical heating system is Y6-VFP (Grant Instruments Ltd., U.K.). The temperature was measured with a mercury-in-glass thermometer. A device consisting of a 2200-type pressure sensor and PDRC-1C/2C type display supplied by the MKS Corporation (Andover MA, USA) was used to measure the pressure directly. The uncertainty of the measured pressure was 0.013 kPa. This apparatus was tested by measuring the vapor pressure of benzyltributylammonium chloride aqueous solutions at different concentrations from (298.15 to 303.15) K as given in Table 2. The average absolute deviations of the experimental data sets were less than 0.02 kPa compared to the published data.³ This test showed the reliability of this apparatus and experimental procedure.

Results and Discussion

The vapor pressures of the binary systems 1,4-dioxane + TBAN and water + TBAN have been measured at (288.15, 298.15, and 308.15) K. In these systems for each temperature, 6 or 11 vapor pressure measurements of different salt concentrations have been studied. The molalities of TBAN ranged from (0.100 to 4.000) $\text{mol} \cdot \text{kg}^{-1}$. The vapor pressures of the 1,4-dioxane + water + TBAN system have also been measured at 11 constant salt molalities (0.100, 0.250, 0.500, 0.750, 1.000, 1.500, 2.000, 2.500, 3.000, 3.500, and 4.000) $\text{mol} \cdot \text{kg}^{-1}$ at

**Figure 2.** Vapor pressure p of 1,4-dioxane in the 1,4-dioxane + tetrabutylammonium nitrate (2) system: Δ , 288.15 K; $\diamond$, 298.15 K; $+$, 308.15 K. m_2 is the salt molality.**Figure 3.** Vapor pressure p of water in the water (1) + tetrabutylammonium nitrate (2) system: $*$, 288.15 K; $\diamond$, 298.15 K; $+$, 308.15 K. m_2 is the salt molality.

(288.15, 298.15, and 308.15) K. All the experimental data given in Tables 3 to 5 and in Figure 2 and Figure 3. The vapor pressures of 1,4-dioxane and water decrease with an increase of salt molality in the binary systems.

Calculation of the Activity Coefficients of Solvents in Binary Systems. Vapor–liquid equilibrium data for the systems 1,4-dioxane (1) + TBAN (2) and water (1) + TBAN (2) were correlated by using the electrolyte NRTL model of Chen et al.¹¹ This model is usually used for correlation and prediction of phase equilibria of the binary electrolyte + non-electrolyte systems. The vapor phase was assumed as an ideal gas. In the case of salt-containing binary systems, the salt was regarded as a nonvolatile component ($y_2 = 0$ and $y_1 = 1$). The formula for the vapor–liquid equilibrium is

$$y_i P = \gamma_i x_i p_i^0$$

The saturation vapor pressure of pure solvent i (p_i^0) at temperature T was calculated with the published Antoine constants.¹³ x_i is the liquid-phase mole fraction of solvent i on the assumption of total dissociation of salt.

The parameters of the electrolyte NRTL model were obtained from the published data¹⁴ and are shown in Table 6. This model was often used to calculate the activity coefficients of solvents γ_i in the binary solvent + electrolyte systems. The activity coefficients of solvents of the studied binary systems are presented in Table 3 and Table 4. The expressions for calculation of the activity coefficients of solvents were directly taken from the electrolyte NRTL model of Chen et al.¹¹

Table 5. Total Vapor Pressure p of the 1,4-Dioxane (1) + Water (2) + Tetrabutylammonium Nitrate (3) System from $T = (288.15 \text{ to } 308.15) \text{ K}$ as a Function of Liquid Mole Fraction of 1,4-Dioxane x'_1 on a Salt-Free Basis and Salt Molality m_3

$m_3/(\text{mol} \cdot \text{kg}^{-1})$	p/kPa			$m_3/(\text{mol} \cdot \text{kg}^{-1})$	p/kPa		
	$T/\text{K} = 288.15$	$T/\text{K} = 298.15$	$T/\text{K} = 308.15$		$T/\text{K} = 288.15$	$T/\text{K} = 298.15$	$T/\text{K} = 308.15$
$x'_1 = 0.050$							
0.000	2.352	4.157	7.238	0.000	3.657	6.357	10.735
0.100	2.369	4.197	7.215	0.100	3.640	6.342	10.684
0.250	2.389	4.194	7.186	0.250	3.596	6.234	10.589
0.500	2.390	4.150	7.126	0.500	3.529	6.135	10.388
0.750	2.381	4.114	7.003	0.750	3.490	5.993	10.190
1.000	2.352	4.090	6.947	1.000	3.437	5.876	10.068
1.500	2.309	4.053	6.839	1.500	3.344	5.682	9.824
2.000	2.264	4.014	6.787	2.000	3.257	5.482	9.645
2.500	2.216	3.966	6.743	2.500	3.196	5.336	9.486
3.000	2.172	3.922	6.696	3.000	3.122	5.192	9.363
3.500	2.136	3.884	6.654	3.500	3.041	5.094	9.227
4.000	2.108	3.846	6.607	4.000	2.968	5.002	9.091
$x'_1 = 0.100$							
0.000	2.709	4.930	8.617	0.000	3.749	6.359	10.828
0.100	2.724	4.965	8.593	0.100	3.718	6.342	10.774
0.250	2.736	4.949	8.542	0.250	3.669	6.257	10.696
0.500	2.730	4.890	8.461	0.500	3.610	6.181	10.543
0.750	2.694	4.828	8.358	0.750	3.562	6.069	10.370
1.000	2.660	4.742	8.245	1.000	3.521	5.965	10.263
1.500	2.576	4.590	8.002	1.500	3.421	5.773	10.042
2.000	2.514	4.441	7.842	2.000	3.338	5.637	9.839
2.500	2.452	4.336	7.663	2.500	3.254	5.542	9.679
3.000	2.386	4.233	7.511	3.000	3.201	5.464	9.541
3.500	2.317	4.141	7.366	3.500	3.158	5.400	9.430
4.000	2.260	4.078	7.216	4.000	3.088	5.336	9.331
$x'_1 = 0.170$							
0.000	2.984	5.482	9.703	0.000	3.386	6.027	9.914
0.100	2.992	5.481	9.661	0.100	3.374	6.001	9.854
0.250	2.980	5.441	9.563	0.250	3.357	5.997	9.823
0.500	2.925	5.348	9.447	0.500	3.332	5.945	9.745
0.750	2.868	5.290	9.349	0.750	3.309	5.862	9.691
1.000	2.814	5.208	9.258	1.000	3.294	5.801	9.626
1.500	2.686	5.032	9.133	1.500	3.264	5.681	9.469
2.000	2.620	4.882	9.007	2.000	3.216	5.549	9.335
2.500	2.537	4.728	8.849	2.500	3.162	5.456	9.155
3.000	2.456	4.585	8.694	3.000	3.108	5.352	9.042
3.500	2.349	4.442	8.509	3.500	3.066	5.285	8.925
4.000	2.300	4.336	8.351	4.000	3.032	5.246	8.806

Table 6. Energy Parameters Δg_{ij} and Δg_{ji} and Nonrandomness Factors α_{ij} for the Electrolyte NRTL Model

component			Δg_{ij}	Δg_{ji}
i	j	α_{ij}	$\text{J} \cdot \text{mol}^{-1}$	$\text{J} \cdot \text{mol}^{-1}$
water	TBAN	0.2	2613.158	-134.126
1,4-dioxane	TBAN	0.1	3369.516	-822.603

The activity coefficients of solvents in the ternary system 1,4-dioxane + water + TBAN were earlier¹⁴ calculated by using the electrolyte NRTL model of Mock et al.¹⁵ which is used for the ternary mixed-solvent electrolyte systems.

Conclusion

The vapor pressures of the 1,4-dioxane + tetrabutylammonium nitrate, water + tetrabutylammonium nitrate, and 1,4-dioxane + water + tetrabutylammonium nitrate systems have been measured at different salt molalities at (288.15, 298.15, and 308.15) K with a modified Othmer-type equilibrium cell. The activity coefficients of solvents in the binary systems were calculated by using the electrolyte NRTL model.

Literature Cited

Mischenko, K. P.; Poltoratskiy, G. M. *Thermodynamics and Structure of Aqueous and Nonaqueous Electrolyte Solutions*, 2nd ed.; Khimiya: Leningrad, 1976; (in Russian).
Belveze, L. S.; Brennecke, J. F.; Stadtherr, M. A. Modeling of Activity Coefficients of Aqueous Solutions of Quaternary Ammonium Salts

with the Electrolyte-NRTL Equation. *Ind. Eng. Chem. Res.* **2004**, *43*, 815–825, and references cited therein.

- Lee, L.-S.; Lee, C.-C. Vapor Pressures and Enthalpies of Vaporization of Aqueous Solutions of Benzyltrimethylammonium Chloride, Benzyltriethylammonium Chloride, and Benzyltributylammonium Chloride. *J. Chem. Eng. Data* **1998**, *43*, 17–20.
- Lee, L.-S.; Lee, C.-C. Vapor Pressures and Enthalpies of Vaporization of Aqueous Solutions of Triethylammonium Chloride, 2-Hydroxyethylammonium Chloride, and Tris(hydroxymethyl)aminomethane Hydrochloride. *J. Chem. Eng. Data* **1998**, *43*, 469–472.
- Lee, L.-S.; Huang, M.-Y.; Hsu, H.-L. Vapor Pressure of Ethanol + Benzyltributylammonium Chloride Solution and Vapor–Liquid Equilibrium of Ethanol + Water + Benzyltributylammonium Chloride Mixture at Atmospheric Pressure. *J. Chem. Eng. Data* **1999**, *44*, 528–531.
- Barthel, J.; Lauermaun, G.; Neueder, R. Vapor Pressure Measurements on Non-aqueous Electrolyte Solutions. Part 2. Tetraalkylammonium Salts in Methanol. Activity Coefficients of Various 1–1 Electrolytes at High Concentrations. *J. Solution Chem.* **1986**, *15*, 851–867.
- Lauermaun, G.; Kunz, W.; Barthel, J. Vapor Pressure Measurements on Nonaqueous Solutions. Part IV. HNC Calculations Using Friedman's Cosphere Overlap Model. *J. Solution Chem.* **1987**, *16*, 871–884.
- Barthel, J.; Kunz, W. Vapor Pressure Data for Non-aqueous Electrolyte Solutions. Part 5. Tetraalkylammonium Salts in Acetonitrile. *J. Solution Chem.* **1988**, *17*, 399–415.
- Barthel, J.; Klein, L.; Kunz, W.; Calmettes, P.; Turq, P. Tetraalkylammonium Bromides in Methanol: Small Angle Neutron Scattering and Vapor Pressure Measurements. *J. Solution Chem.* **1994**, *23*, 955–971.
- Barthel, J.; Neueder, R.; Wittmann, H. Osmotic Coefficients and Activity Coefficients of Nonaqueous Electrolyte Solutions. Part 3.

- Tetraalkylammonium Bromides in Ethanol and 2-Propanol. *J. Solution Chem.* **1999**, 28, 1263–1276.
- (11) Chen, C.-C.; Britt, H. I.; Boston, J. F.; Evans, L. B. Local Composition Model for Excess Gibbs Energy of Electrolyte Systems. Part I: Single Solvent, Single Completely Dissociated Electrolyte Systems. *AIChE J.* **1982**, 28, 588–596.
- (12) Yang, S.-O.; Lee, C.-S. Vapor–Liquid Equilibria of Water + Methanol in the Presence of Mixed Salts. *J. Chem. Eng. Data* **1998**, 43, 558–561.
- (13) Walas, S. M. *Phase Equilibria in Chemical Engineering*; Butterworth: London, 1985.
- (14) Kurzin, A. V.; Evdokimov, A. N.; Poltoratskiy, G. M.; Platonov, A. Y.; Gusev, V. E.; Golubeva, Y. M. Isothermal Vapor–Liquid Equilibrium Data for the Systems 1,4-Dioxane + Water + Tetrabutylammonium Nitrate and Acetonitrile + Water + Tetrabutylammonium Bromide. *J. Chem. Eng. Data* **2004**, 49, 208–211.
- (15) Mock, B.; Evans, L. B.; Chen, C.-C. Thermodynamic Representation of Phase Equilibria of Mixed-Solvent Electrolyte Systems. *AIChE J.* **1986**, 32, 1655–1664.
- (16) Boublic, T. *The Vapor Pressures of Pure Substances*; Elsevier: Amsterdam, 1984.
- (17) Rabinovich, V. A.; Khavin, Z. Ya. *Brief Chemical Handbook*, 4th ed.; Khimiya: Leningrad, 1991; (in Russian).

Received for review September 6, 2007. Accepted October 19, 2007.
V.B.A. thanks the Government of Saint-Petersburg for the financial support [Grant for students M06-3.6D-14].

JE700512A